

EXPRESS MAIL LABEL NO.:

EL804518659US

DATE: November 15, 2001

FORM PTO 1390 (REV 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER ICON-001	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 09/980542	
INTERNATIONAL APPLICATION NO. PCT/US00/13555		INTERNATIONAL FILING DATES 17 May 2000		PRIORITY DATE CLAIMED 17 May 1999	
TITLE OF INVENTION PROCESS OF RAPID VARIETY-INDEPENDENT PLANT TRANSFORMATION					
APPLICANT(S) FOR DO/EO/US Nikolay V. Kuchuk and Victor Klimyuk					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<p>1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing 35 U.S.C. 371</p> <p>3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371 (f)). The submission must include items (5), (6), (9) and (21) indicated below.</p> <p>4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c)(2))</p> <p>a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).</p> <p>b. <input checked="" type="checkbox"/> has been communicated by the International Bureau.</p> <p>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371 (c)(2)).</p> <p>a. <input type="checkbox"/> is attached hereto.</p> <p>b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4).</p> <p>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))</p> <p>a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</p> <p>b. <input type="checkbox"/> have been communicated by the International Bureau.</p> <p>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p>d. <input checked="" type="checkbox"/> have not been made and will not be made.</p> <p>8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).</p> <p>9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). (Unexecuted)</p> <p>10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).</p> <p>Items 11 to 20 below concern document(s) or information included:</p> <p>11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. w/PTO-1449, 5 references</p> <p>12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</p> <p>13. <input type="checkbox"/> A FIRST preliminary amendment.</p> <p>14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.</p> <p>15. <input type="checkbox"/> A substitute specification.</p> <p>16. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.</p> <p>18. <input checked="" type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).</p> <p>19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).</p> <p>20. <input checked="" type="checkbox"/> Other items or information: Copy Of International Preliminary Examination Report, Six (6) Sheets of Formal Drawings</p>					

U.S. APPLICATION NO. (known use 37 CFR 1.5) <div style="font-size: 2em; font-weight: bold; margin-left: 100px;">09/980542</div>	INTERNATIONAL APPLICATION NO. PCT/US00/13555	ATTORNEY'S DOCKET NUMBER ICON-001
--	---	--------------------------------------

21. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00

☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00

☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00

☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00

☒ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

Surcharge of \$ _____ for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	39-20 =	19	x 18.00
Independent claims	2-3 =		x
MULTIPLE DEPENDENT CLAIM(s) (if applicable)			+ 280.00
TOTAL OF ABOVE CALCULATIONS =			
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.			
SUBTOTAL =			
Processing fee of \$ _____ for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).			+
TOTAL NATIONAL FEE =			
Fee for recording the enclosed assignment (37 CFR 1.21 (h)). Assignment must be accompanied by appropriate cover sheet (37 CFR 3.28, 3.31) (_____ per property).			+
TOTAL FEES ENCLOSED =			

Amount to be Refunded:	\$
Charged:	\$

CALCULATIONS PTO USE ONLY

a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 12-1095 in the amount of \$ 361.00 to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required or credit any overpayment to my Deposit Account No. 12-1095. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

LERNER, DAVID, LITTENBERG, KRUMHOLZ & MENTLIK, LLP

600 South Avenue West
Westfield, New Jersey 07090
(908) 518-6346

Shawn P. Foley

SIGNATURE: _____

NAME: Shawn P. Foley

REGISTRATION NUMBER: 33,071

EXPRESS MAIL LABEL NO.: EL804519019US
DATE: March 4, 2002

09/980542

Rec'd PGT/PTO 04 MAR 2002

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of	:	
Nikolay V. Kuchuk, et al.	:	
International Application No. PCT/US00/13555	:	Group Art Unit:
International Filing Date: May 17, 2000	:	Examiner:
For: PROCESS OF RAPID VARIETY-INDEPENDENT	:	Date: March 4, 2002
PLANT TRANSFORMATION	:	
	X	

Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Preliminary to initiation of the prosecution of the above-identified pending U.S. patent application, the following amendment and remarks are respectfully submitted.

IN THE SPECIFICATION

Page 1, following the title, please insert the following heading and paragraph:

CROSS REFERENCE TO RELATED APPLICATIONS

The present application claims benefit of United States Provisional Patent Application 60/134,459, filed May 17, 1999. The present application is also a §371 of International Application PCT/US00/13555, filed May 17, 2000. The disclosures of all of said applications are incorporated by reference herein.

REMARKS

The specification has been amended to include a cross-reference to the prior application.

Applicant respectfully requests that this amendment be entered, and that prosecution on the merits of this application be initiated. If, however, for any reason the

09/980542 "030402"

Examiner does not believe such action can be taken, it is respectfully requested that he telephone applicant's attorney at (908) 654-5000 in order to overcome any objections which he may have.

If there are any additional charges in connection with this requested amendment, the Examiner is authorized to charge applicant's Deposit Account No. 12-1095 therefor.

Respectfully submitted,

LERNER, DAVID, LITTENBERG,
KRUMHOLZ & MENTLIK, LLP


SHAWN P. FOLEY
Reg. No. 33,071

600 South Avenue West
Westfield, NJ 07090-1497
Telephone: (908) 654-5000
Facsimile: (908) 654-7866

09980542 030400

PROCESS OF RAPID VARIETY-INDEPENDENT PLANT TRANSFORMATION

5 FIELD OF THE INVENTION

The present invention relates to methods of introducing genetic material of interest into plants, and more particularly to methods involving transformation and line conversion of plant species that have proven difficult to manipulate on a genetic level.

BACKGROUND OF THE INVENTION

10 Methodologies have evolved during the last twenty years to genetically engineer plants. In general, they are based on either direct DNA introduction into plant cells or indirect transfer mediated by *Agrobacterium tumefaciens*. Methods involving direct transfer include particle bombardment of cultured plant tissues and DNA introduction into
15 naked plant cells *i.e.*, protoplasts, using polyethylene glycol or electroporation. See, *e.g.*, Sawahel & Cove, *Biotech. Adv.* 10:394-412 (1992); Christou, *Cur. Opinion Biotech.* 4:135-141 (1993); Gelvin, *Cur. Opinion Biotech.* 9:227-232 (1998) and Birch, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48:297-326 (1997) Most methods are variety-specific because they are based on use of *in vitro* grown regenerable plant systems which in turn are variety-specific. Except for a few economically important crops such as potato, tomato and canola,
20 transformation methods available currently work with only a handful of varieties.

The traditional backcross method of breeding has provided a mechanism for the transfer of a trait from one line (the donor) to another line (the recurrent parent). See, *e.g.*, Harlan and Pope, *J. Heredity*, 13:319-322 (1922). It has been particularly useful for corn, soybean and cotton. Successful backcross breeding requires: a previously derived
25 recurrent parent; maintenance of the trait of interest during selection; sufficient backcrosses to reconstitute the genome of the recurrent parent. Allard, *Principles of Plant Breeding*, Wiley and Sons (1960). During the backcross program, the hybrid population becomes increasingly homozygous for genes of the recurrent parent at a rate described by the formula:

$$\text{Proportion of homozygosity} = 1 - 0.5^m$$

30 where *m* is the number of backcross generations. Using this formula, one can calculate that more than 98% of the hybrid genome will be homozygous for genes of the recurrent parent after six generations. The formula, however, only describes regions of the genome that are unlinked with respect to the genes being introgressed. The rate at which linked regions approach homozygosity is dependent upon the chromosome recombination frequency. In one

of the most detailed studies assessing the effectiveness of traditional backcross breeding, eight Tm-2-converted isogenic lines of tomato were examined at nine flanking restriction fragment length polymorphism (RFLP) loci. See, Young and Tanksley, *Theor. Appl. Genet.* 77:353-359 (1989). The minimum donor chromosome fragment found after 10 generations of backcrossing and maintained without reduction in size for an additional nine generations was 4 cM the maximum size found even after 11 generations was 51 cM (*i.e.*, more than half of the corresponding chromosome). In marker-assisted selection based on simple sequence repeats (SSR) or RFLP, the reconstruction could be done faster and cleaner, but it would require screening of sizable populations of progeny using relatively expensive methods and would be complicated by the random insertion of transgenes in independent primary transformants.

Plainly, backcrossing is not a trivial task because for most crop plants, hundreds of lines, hybrids or varieties are needed simultaneously. In 1998 for example, the U.S. soybean seed market consisted of over 500 varieties, and the U.S. maize seed market included over 600 hybrids. Another major disadvantage of the backcrossing method is that it is very time- consuming. Line conversion through recurrent backcrossing normally requires 3 to 5 subsequent backcrosses, thus adding at least two and sometimes up to four years to variety development time.

There are many disadvantages associated with current transformation methods. Line conversion, for example, is a process whereby heterologous DNA is transferred from one plant species or variety to another using various forms of sexual (*i.e.*, pollination) or somatic (*i.e.*, cellular) hybridization. Because current methods are species- and variety-specific, they can be commercially used only in combination with line conversion technologies that allow for transgene transfer from a primary transformant into multiple varieties of interest. In addition, they result in random transgene insertion into the host genome. Therefore, extensive screening of numerous independent transformation events are required in order to identify the events that are stable, inheritable and allow for proper transgene expression. Subsequent transgene insertions cannot be addressed to the same site, thus complicating breeding. Linkage drag (*i.e.*, co-inheritance of unwanted traits) and a limited ability to handle multiple independent transgene traits present even further difficulties.

SUMMARY OF THE INVENTION

Applicants have invented a method of introducing nucleic acid into plants and producing genetically engineered plants. The methods are applicable to plants such as corn and wheat which have been quite difficult to genetically modify by existing technologies. In addition, the method constitutes a significant improvement over backcrossing methods because it achieves the same or better results in a much shorter time period.

The nucleic acid of interest is not introduced directly into the plant of interest or what is referred to as the recipient. Rather, it is first routed to another plant, different from the recipient, and which is referred to as the donor or the clipboard species. The nucleic acid is then moved from the donor to the recipient. One method entails sexual hybridization or "crossing" the two plants. Pollen from the donor is used to pollinate a recipient plant. Another method is conducted on the cellular level whereby cells or protoplasts of the donor and the recipient plants are fused. One feature of the invention allows for the nucleic acid to be moved from the donor to the recipient without the movement of any native genomic DNA of the donor. This is accomplished due to the selection of donor/recipient pairs that normally produce unstable hybrids. This means that the respective genomes are unstable and thus do not commingle so as to produce a "hybrid" plant. This phenomenon is hereinafter referred to as "producing unstable progeny or demonstrating preferential segregation or sorting out." During the temporary coexistence of the chromosomes of the donor and the recipient, the nucleic acid or the gene of interest is moved to the genome of the recipient plant. Another feature of the present invention accomplishes random or site-specific introduction of nucleic acid by surrounding the nucleic acid of interest with flanking sequences that allow transposition of the nucleic acid into a random location or direct the insertion of the nucleic acid into a specific location in the genome of the recipient.

Accordingly, one aspect of the present invention is directed to a method for introducing genetic material into plants, comprising:

preparing a first plant transformed with a heterologous nucleic acid having 5' and 3' excisable flanking sequences that allow movement of the heterologous nucleic acid from one genome to another;

crossing a second plant and the transformed first plant, wherein the first and second plants, upon crossing, produce unstable progeny or demonstrate preferential segregation or sorting out; and

selecting progeny of the second plant of (b) which contain the heterologous nucleic acid.

In preferred embodiments, the 5' and 3' excisable flanking sequences comprise a transposable element, and the first plant, the second plant or both the first plant and the second plant produce a transposase specific to the transposable element. In another preferred embodiment, the 5' and 3' excisable flanking sequences are recombination sites and the first plant, the second plant or both the first and second plants produce a recombinase specific to the recombination sites:

In other preferred embodiments, the first plant, also referred to as the donor or the clipboard species, is *Tripsacum* and in the second plant, also referred to as the recipient, is maize, wheat, barley or oat. In another preferred embodiment, the donor is *Orychophragmus* and the recipient is a crucifer such as canola. Other preferred donor/recipient pairs are: *Glycine tomentella*/soybean, *Solanum phreja*/potato, maize/wheat, maize/barley, maize/oat, *Pennisetum*/wheat, *Pennisetum*/barley, *Hordeum bulbosum*/barley, *Hordeum bulbosum*/wheat, *Nicotiana digluta*/*Nicotiana tabacum* and *Oryza minuta*/rice.

In other preferred embodiments, the donor and/or the recipient plant carries a *Se* semigamy mutation. In yet other preferred embodiments, the donor and/or recipient plant is soybean carrying a *ms* mutation causing polyembryony.

In other preferred embodiments of the process, transgenes are targeted into specific predefined genome sites through targeted recombination as an integrative locus.

A related aspect of the present invention is directed to a method for introducing genetic material into plants that is conducted on a somatic level. This method involves the following steps:

preparing a cell or protoplast of a first plant transformed with a heterologous nucleic acid having 5' and 3' excisable flanking sequences that allow movement of the heterologous nucleic acid from one genome to another;

fusing the cell or protoplast with a cell or protoplast of a second plant to produce a fused cell or a fused protoplast, wherein the first and second plants, upon crossing, produce unstable progeny or demonstrate preferential segregation or sorting out;

regenerating whole plants from the fused cell or the fused protoplast; and

selecting progeny of the regenerated plants that contain the heterologous nucleic acid. The fused cells or protoplasts *per se*, are also provided. Further, the methods of the present invention produce plants that have a different genetic make-up than transgenic

plants made by other methods because the end result of the process is an individual plant that is genetically devoid of any resident DNA of primary transformant (*i.e.*, the donor). Progeny of the plant, plant parts and seed and seed parts from the plant are also provided.

The methods of the present invention provide for transgene manipulation in essentially all crop species, especially the economically important varieties. The methods are not only generally applicable to essentially all crop species, but they are rapid (one to two crosses), free of linkage drag and variety-independent. In addition, the methods described are the only variety-independent process of transformation and line conversion that can be used for genetic engineering of complex lines/hybrids which cannot be recovered after crosses with other varieties.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a linear plasmid map of pIC156;

Fig. 2 is a linear plasmid map of pIC216;

Fig. 3 is a linear plasmid map of pIC312;

Fig. 4 is a linear plasmid map of pIC31A2;

Fig. 5 is a linear plasmid map of pIC401; and

Fig. 6 is a linear plasmid map of pIC411.

DETAILED DESCRIPTION OF THE INVENTION

The methods of the present invention produce transformed plants by transforming a donor species or mutant with a construct capable of excision/reinsertion, preferably by a transposon-mediated or homologous or non-homologous recombination mechanism, crossing the donor with a recipient, whereas donor and recipient organisms have been selected from species/mutant combinations that upon sexual/somatic hybridization produce hybrids that are unstable and demonstrate genome instability and segregation of one or both pure parental genotypes, inducing or selecting for excision of the heterologous nucleic acid from the recipient and integration into the donor parental chromosome, and lastly, selecting a progeny that is essentially a genetically pure recipient plant that carries the transgene in question. The flow of heterologous genetic material is completely separated from resident gene flow during genetic manipulations by utilizing species or mutant combinations of recipient and donor organisms that upon sexual/somatic hybridization produce hybrids that show no recombination between homologous/homologous chromosomes and which are unstable and upon mitotic or meiotic divisions, sort out pure parental genomes of one or both types.

By the term "plant", it is meant to include all flowering plants, and all forms, lines and varieties of the plant. "Transformed" is used herein to mean genetically modified by the incorporation of heterologous DNA into cells. By the term "heterologous", it is meant DNA not normally found in the recipient plant.

Species-specific chromosome elimination (genome segregation) in interspecific/intergeneric hybrids is a well-documented phenomenon. In many cases, however, unstable hybrids were of limited interest as the main breeding efforts were aimed at chromosome exchange between two parental genomes as a method for introgression of alien chromosomal material. Prior to the time the present invention was made, unstable hybrids segregating parental genomes were described only in terms of systems that produce haploid plants (interspecific, intergeneric crosses for production of haploid wheat, barley, potato) or in terms of negative results of attempts to achieve an introgression of chromosome material from wild species into cultivated crops (such as from *Tripsacum* to maize or *Glycine tomentella* to soybean).

In general, for every crop species (including all varieties and lines thereof), there is a wild relative or a mutant form, that, upon hybridization, forms an unstable hybrid and can serve as a donor or clipboard plant as defined herein. An empirical way to identify such an organism involves crossing a crop species of interest with a number of related species and testing the genetic makeup of the resulting progeny. Methods of preliminary identification of progeny that is predominantly uniparental are known in the literature and are based on different selective or non-selective traits. Methods of broad and reliable genotyping of the progeny are numerous, simple, and rely on analysis of various markers in genomic DNA. Based on such primary screening and subsequent genotyping, suitable clipboard organisms can be rapidly identified. Beyond this basic criterion, the donor/recipient pairs are chosen so as to provide a an adequate duration of the hybrid state in cells of primary hybrid, or of its progeny. While complete elimination is a desired end state, relative duration of coexistence of chromosomes of both species in the same cell is important as it provides sufficient time for the excision of transgene locus from the "clipboard" organism and its integration into the chromosome of recipient's chromosome.

Physical interaction and in particular, chromosome material exchange between parental genomes in a hybrid, can be excluded or minimized in a number of different ways. The most well-known approach is based on the use of interspecific or intergeneric hybridization -- the hybrids produced show little if any homologous chromosome pairing,

thus limiting crossover. In addition, many of such hybrids are more or less genetically unstable and show tendency for a rapid elimination of one parental genome. As a result, by using such a cross, the chromosomes of two remote parental organisms may be placed in a hybrid nucleus to produce a hybrid state for directed exchange of transgene material between the parents. However, resident chromosome material of the two parents essentially does not interact, and subsequent chromosome elimination will allow for elimination of one parent as soon as F₀, F₁ or BC₁ progeny.

In addition to hybridization between remote species as a process that allows for a temporary hybrid state followed by rapid recovery of pure parental genomes, there are other approaches that achieve similar results. One such approach is based on the use of mutants that reduce or eliminate chromosome crossover and/or those that cause pure parental genome segregation in intra-specific as well as interspecific crosses. One example is semigamy in cotton, a mutation which causes sperm nucleus to enter egg cell but subsequent nuclear fusion supposedly does not take place and both nuclei divide independently resulting in F₁ plants that are chimeral for sectors of haploid tissues of paternal and maternal type. See, Turcotte and Feaster, *J. Hered.* 58:55-57 (1967). Another approach is the well-characterized *Oenothera* system in which all chromosomes are involved in translocations in such a way that the F₁ crosses with normal stocks will have at meiosis a ring containing the entire haploid number of chromosomes, thus excluding independent chromosome sorting out. A further approach to the use of chemical/physical treatments such as irradiation (Pandey, *N. Z. J. Bot.*, 18:203-207 (1980)) or other, of one parent that result in a damage and subsequent preferential elimination of damaged genome. Such an approach has been instrumental in production of gynogenetic onion plants after pollinating with gamma-irradiated pollen. See, Dore & Marie, *Plant Breeding* 111:142-147 (1993).

To carry out the present invention for any given plant, a wild or remote relative can be found that allows for genetically unstable hybrids that are characterized by rapid genome segregation. Well-characterized combinations involving economically important crops are summarized below.

Insofar as dicotyledonous crops are concerned, the best studied case for potato is a hybridization between commercial varieties of potato, *Solanum tuberosum*, and a wild species *Solanum phureja*, that results in high frequency of haploid production as a result of early *phureja* chromosome elimination in hybrid embryo - Hougas, *et al.*, *Crop. Sci.* 4:593-

595 (1964); Clulow, *et al.*, *Theor. Appl. Genet.* 82:545-551 (1991). Regarding Canola/Rapeseed, somatic separation of the parental genomes in hybrids between *Brassica napus* and *Orychophragmus violaceus* is described in Li, Z., *et al.*, *Theor. Appl. Genet.* 91:131-136 (1995); Li, *et al.*, *Hereditas* 125: 69-75 (1996); Li, *et al.*, *Theor. Appl. Genet.* 96:251-265 (1998); Wu, J., *et al.*, *Plant Breeding* 116:251-257 (1997). The hybrid is morphologically intermediate but is self-fertile and upon selfing produces mostly *B. napus* progeny. *Orychophragmus* method works also with *Brassica juncea* and *Brassica carinata*, two other *Brassica* species of economic importance. It will also operate with other economically crucifers such as *B. oleracea*, *B. campestris*, *Raphanus sativus*. Regarding soybean, wild species genome elimination in the progeny of a hybrid between soybean and *Glycine tomentella* is documented in Shoemaker, *et al.*, *Theor. Appl. Genet.* 80:17-23 (1990). Yet other examples of matroclinous plants result from crosses between *Fragaria vesca* (strawberries) and *Fragaria chiloensis* or *F. virginiana* (Ichijima, *Genetics*, 11:590-604 (1926)) as well as patroclinous plants from crosses between *Nicotiana glauca* and *N. tabacum* (tobacco, Clausen and Lammerts, *Amer. Nat.* 63:279-322 (1929)).

Insofar as monocotyledonous crops are concerned, Galinat, *Ann. Rev. Genet.*, 5: 447-478 (1971) and Galinat, *Evolution*, 27:644-655 (1973), demonstrated that in a cross between a diploid maize and a diploid *Tripsacum dactyloides*, the F₁ hybrid had the expected amphihaploid chromosome number. *Tripsacum* chromosomes could not pair in meiosis, however, and since *Tripsacum* chromosomes tend to be lost during mitosis as well as meiosis, diploid maize was recovered as soon as BC₁. These results have been routinely reproduced in numerous breeding labs around the world. When wheat or barley is crossed with wild species *Hordeum bulbosum*, a high frequency of haploids is obtained as a result of fertilization and subsequent elimination of *bulbosum* genome (Kasha and Kao, *Nature* 225:874-876 (1970); Barclay, *Nature* 256:410-411 (1975)). The method has been widely used for haploid production of numerous varieties of both crops. *Bulbosum* method has been replaced by wide crosses, wherein wheat (both *Triticum aestivum* as well as *T. turgidum*), triticale or barley plants are pollinated by pollen of maize, sorghum, pearl millet or *Tripsacum*. The resultant hybrids are highly unstable and as a rule, developing plants retain only maternal genome. This haploidy method works with dozens of wheat varieties and is essentially variety-independent. Laurie and Bennett, *Theor. Appl. Genet.* 76:393-397 (1988); Ohkawa, *et al.*, *Jap. J. Breed.* 42:891-894, (1992); Ushiyama, *et al.*, *Jap. J. Breed.* 41:353-357 (1991); Furusho, *et al.*, *Jap. J. Breed.* 41:175-179 (1991); Laurie, *Genome*, 32:1063-1067

(1989). The same method is also applicable for haploid production in oats. See, Rines and Dahleen, *Crop Sci.*, 30, 1073 (1990). Preferential genome segregation also occurs in progenies of interspecific rice (*Oryza*) combinations, such as *O. sativa* and *O. minuta*. See, Mariam, *et al.*, *Theor. Appl. Genet.* 93:664-671 (1996).

5 In summary, the literature provides examples of numerous, well characterized hybridization combinations that when used in accordance with the present invention, allow for the creation of a temporary hybrid state during which heterologous genetic material may be exchanged. In addition, many wild species may be used as transgene donors for rapid and linkage drag-free transgene transfer into multiple species of important crops. Wild species
10 include *Tripsacum* (e.g., for maize, wheat, barley and oats); *Oryza minuta* (e.g., for rice), *Orychophragmus* (e.g., for canola and other economically important crucifers); *Solanum phureja* (e.g., for potato), and *Glycine tomentella* (e.g., for soybean). Mutants such as a semigamy mutant of cotton or a ms mutation causing polyembryony in soybean may also be used for the same purpose. Similar species combinations or mutants may be easily identified
15 for other important crop species including sugar beets, peas and tomatoes. The donor species is transformed with an exogenous or heterologous construct that contains genetic information, the genetic instructions necessary to direct the excision of the transgene in question and its reintegration into another chromosome based on either homologous or non-homologous recombination mechanisms, and the DNA of interest. To facilitate selection of successful
20 transformants, the construct also contains DNA encoding a selectable marker such as a trait to be monitored (e.g., antibiotic or herbicide resistance or a phenotypic marker, in particular beta-glucuronidase and/or green fluorescent protein). The DNA of interest may contain one or more genes encoding different proteins. Expression of the genes in progeny of the recipient plant may result in greater resistance to fungal, viral and/or bacterial diseases, pests,
25 insecticides or environmental stress, or may result in improved flavor, storage or nutritional properties. The recipient organism may be an untransformed plant or a plant transformed to contain in its genome specific sites necessary for homologous recombination exchange with the exogenous DNA inserts in donor genome.

30 In preferred embodiments, the heterologous DNA is moved from the donor to the recipient via a system based on (1) transposon-mediated non-homologous transgene transfer or (2) targeted transfer utilizing homologous recombination mechanisms. Transposons are mobile genetic elements that can comprise a substantial part of the genome of a plant and create tremendous phenotypic diversity. Transposable elements are mobile

segments of DNA capable of excision and reinsertion into another locus on a chromosome. Plant transposons are among first mobile DNA elements described and a number of plant transposable elements that have been cloned, such as Ac/DS, Mu and En/Spm, are preferred for use in the present invention. These transposable elements are currently used as genetic tools in plant molecular biology and biotechnology. They serve as invaluable tools for plant developmental studies and for plant genome analysis and plant gene isolation through the so-called insertional mutagenesis and tagging. See, e.g., Walbot, *Ann. Rev. Plant Mol. Biol.* 43:49-82 (1992). Other examples of transposable elements for use in the present invention are described in Fedoroff, U.S. Patent 4,732,856; Dooner *et al.*, PCT Application WO91/156074; etc.), Yoder and Lassner, PCT Application WO92/01370, and Ebinuma *et al.*, PCT Application WO96/15252.

For the purposes of the present invention, transposon-based excision and reinsertion of transgenes is a system that allows for the disconnection of the transgene movement from the resident plant gene movement during crosses while providing an additional important advantage of full instructions directing transgene excision and reinsertion in just one genetic construct and via one transformation step. Thus, the system does not require genetic engineering of landing sites in recipient organisms.

In another preferred embodiment, the heterologous DNA is integrated in a specific site of the recipient genome by use of a recombinase and recombination site combination. Site-specific recombinases from bacteriophage and yeasts are being widely used as tools for manipulating DNA both in the test-tube and in living organisms. Preferred recombinases/recombination site combinations for use in the present invention are Cre-*Lox*, FLP-*FRT*, and R-*RS*, where Cre, FLP and R are recombinases, and *Lox*, *FRT*, and *RS* are the recombination sites. Other suitable systems include the intron-encoded yeast endonuclease I-SceI, may be used. See, Choulika, *et al.*, *Mol. Cell Biol.* 15:1968-1973 (1995). To be functional in plants, these sites require 7-8 base pairs (bp) of core sequence between 12-13 bp inverted repeats; the asymmetric core site determines the site orientation, and thus the types of recombination product. Regardless of whether recombination sites are placed on or within a single DNA molecule in direct or opposite orientation, or placed on unlinked linear or circular DNA molecules, the corresponding recombinase can catalyze the reciprocal exchange to produce a deletion, inversion, translocation or co-integration event. See, Bollag, *et al.*, *Ann. Rev. Genet.* 23:199-225 (1989); Kilby, *et al.*, *Trends Genet.* 9:413-421 (1993); and Ow, *Curr. Opinion Biotech.* 7:181-186 (1996). In the present invention, recombinase-

mediated site-specific translocation occurs between different, and in particular non-homologous chromosomes. This in-trans recombinase effect is essential in order to effect transfer of transgenes between two chromosomes belonging to different parents in a hybrid. See, Dale and Ow, *Gene* 91:79-85 (1990); Odell, *et al.*, *Mol. Gen. Genet.* 223:369-378 (1990); Dale and Ow, *Proc. Natl. Acad. Sci. USA* 88:10558-10562 (1991); Russell, *et al.*,
5 *Mol. Gen. Genet.* 234:49-59 (1992); Lyznik, *et al.*, *Plant J.* 8:177-186 (1995); Albert, *et al.*, *Plant J.* 7:649-659 (1995); van Deursen *et al.*, *Proc. Natl. Acad. Sci. USA* 92:7376-7380 (1995).

Examples of suitable homologous recombination systems for use in the present invention are disclosed in the literature, including the Cre-Lox system (Sauer, U.S. Patent 4,959,317, Odell, *et al.*, U.S. Patent 5,658,772; Odell, *et al.*, PCT WO91/09957) and the FLP-FRT system (Hodges and Lyznik, U.S. Patent 5,527,695). One particular utility of known recombination systems for transgene management in plants is directed excision of a transgene from plant genome, a procedure that allows elimination of unwanted heterologous
10 genetic material such as antibiotic selective markers from a commercial variety (Ow and Dale, PCT WO93/01283). These systems, however, address an entirely different utility area, namely, the use of homologous recombination to eliminate unwanted portions of heterologous DNA, rather than to manage separation of flows of transgenes and resident plant genes. Another utility is described in Hooykaas and Mozo, U.S. Patent 5,635,381, and
15 Offringa, *et al.*, U.S. Patent 5,501,967, directed to the use of homologous recombination systems to achieve a site-directed targeted integration of DNA into plant genomes via *Agrobacterium*-mediated transformation. These cases also are limited to targeted transfer between bacteria and plant cells rather than between two plant organisms.
20

Homologous recombination-based transgene shuffling has both clear and strong advantages. By employing precise targeting via homology-addressed DNA sites,
25 transgene "landing sites" can be created that are carefully selected and characterized in advance. As a result, higher level of predictability and reproducibility of transgene behavior, including heritability, expression level, absence of silencing, etc., is achieved. Also, later versions of the transgene cassette can be addressed to the same site, replacing old versions of transgenes with newer ones. Subsequent breeding of the material with a preselected and
30 determined and mapped integration sites is much easier and straightforward. Those skilled in the art will appreciate that this system can be used only if all recipients have been "pre-wired" to contain integration sites. Such an introgression is possible by using other transfer

mechanisms, for example, transposon-mediated transfer or classical introgression by backcrossing. In more specific cases, a recombinase gene also can be introduced into acceptor species in addition to its recombination site. In such cases, the recombinase gene will be under control of artificial transcription factor-mediated promoter where transcription factor is constitutively expressed by the gene located in donor (clipboard) plant. Recombinase can be produced only during the co-existence of two genomes in unstable hybrids. Alternatively, recombinase can be located in clipboard plant, but the transcription factor is constitutively expressed in the recipient plant.

The heterologous DNA may be introduced into the donor plant in accordance with standard techniques. Transformation techniques for dicotyledons are well known in the art and include *Agrobacterium*-based techniques and techniques which do not require *Agrobacterium*. Non-*Agrobacterium* techniques involve the uptake of exogenous genetic material directly by protoplasts or cells. These techniques include PEG or electroporation mediated uptake, particle bombardment-mediated delivery and microinjection. Examples of these techniques are described in Paszkowski *et al.*, EMBO J 3:2717-2722 (1984), Potrykis *et al.*, Mol. Gen. Genet. 199:169-177 (1985), Reich *et al.*, Biotechnology 4:1001-1004 (1986), and Klein *et al.*, Nature 327:70-73 (1987). In each case, the transformed cells are regenerated to whole plants using standard techniques.

Agrobacterium-mediated transformation is a preferred technique for transformation of dicotyledons because of its high efficiency of transformation and its broad utility with many different species. The many crop species which are routinely transformable by *Agrobacterium* include tobacco, tomato, sunflower, cotton, oilseed rape, potato, soybean, alfalfa and poplar (EP 0 317 511 (cotton), EP 0 249 432 (tomato), WO 87/07299 (*Brassica*), U.S. Patent 4,795,855 (poplar)). *Agrobacterium* transformation typically involves the transfer of the binary vector carrying the foreign DNA of interest (e.g., pCIB200 or pCIB2001) to an appropriate *Agrobacterium* strain which may depend on the complement of *vir* genes carried by the host *Agrobacterium* strain either on a co-resident plasmid or chromosomally (e.g., strain CIB542 for pCIB200 (Uknes *et al.*, Plant Cell 5:159-169 (1993))). The transfer of the recombinant binary vector, to *Agrobacterium* is accomplished by a triparental mating procedure using *E. coli* carrying the recombinant binary vector, a helper *E. coli* strain which carries a plasmid such as pRK2013 which is able to mobilize the recombinant binary vector to the target *Agrobacterium* strain. Alternatively, the recombinant

binary vector is transferred to *Agrobacterium* by DNA transformation (Höfgen & Willmitzer, Nucl. Acids Res. 16, 9877 (1988)).

Transformation of the target plant species by recombinant *Agrobacterium* usually involves co-cultivation of the *Agrobacterium* with explants from the plant and follows protocols known in the art. Transformed tissue is regenerated on selectable medium carrying an antibiotic or herbicide resistance marker present between the binary plasmid T-DNA borders.

Preferred transformation techniques for monocots include direct gene transfer into protoplasts using PEG or electroporation techniques and particle bombardment into callus tissue. Transformation can be undertaken with a single DNA species or multiple DNA species (*i.e.* co-transformation) and both these techniques are suitable for use with this invention. Co-transformation may have the advantage of avoiding complex vector construction and of generating transgenic plants with unlinked loci for the gene of interest and the selectable marker, enabling the removal of the selectable marker in subsequent generations, should this be regarded desirable. However, a disadvantage of the use of co-transformation is the less than 100% frequency with which separate DNA species are integrated into the genome (Schocher *et al.*, Biotechnology 4:1093-1096 (1986)).

Published Patent Applications EP O 292 435, EP O 392 225 and WO 93/07278 describe techniques for the preparation of callus and protoplasts of maize, transformation of protoplasts using PEG or electroporation, and the regeneration of maize plants from transformed protoplasts. Gordeon-Kamm, *et al.*, Plant Cell 2:603-618 (1990), and Fromm, *et al.*, Biotechnology 11:194-200 (1993), describe techniques for the transformation of elite inbred lines of maize by particle bombardment.

Transformation of rice can also be undertaken by direct gene transfer techniques utilizing protoplasts or particle bombardment. Protoplast-mediated transformation has been described for *Japonica*-types and *Indica*-types (Zhange, *et al.*, Plant Cell Rep. 7:739-744 (1988); Shimamoto, *et al.*, Nature 338:274-277 (1989); Datta, *et al.*, Biotechnology 8:736-740 (1990)). Both types are also routinely transformable using particle bombardment (Christou, *et al.*, Biotechnology 9:957-962 (1991)).

Patent Application EP 0 332 581 describes techniques for the generation, transformation and regeneration of *Pooideae* protoplasts. Furthermore wheat transformation is described in Vasil, *et al.*, Biotechnology 10:667-674 (1992) using particle bombardment into cells of type C long-term regenerable callus, Vasil, *et al.*, Biotechnology 11:1553-1558

(1993) and Weeks, *et al.*, Plant Physiol. 102:1077-1084 (1993) describe particle bombardment of immature embryos and immature embryo-derived callus.

Transformation of monocot cells such as *Zea mays* is achieved by bringing the monocot cells into contact with a multiplicity of needle-like bodies on which these cells may be impaled, causing a rupture in the cell wall thereby allowing entry of transforming DNA into the cells. See U.S. Patent 5,302,523. Transformation techniques applicable to both monocots and dicots are also disclosed in the following U.S. Patents: 5,240,855 (particle gun); 5,204,253 (cold gas shock accelerated microprojectiles); 5,179,022 (biolistic apparatus); 4,743,548 and 5,114,854 (microinjection); and 5,149,655 5,120,657 (accelerated particle mediated transformation); 5,066,587 (gas driven microprojectile accelerator); 5,015,580 (particle-mediated transformation of soy bean plants); 5,013,660 (laser beam-mediated transformation); 4,849,355 and 4,663,292.

The thus-transformed plant cells or plant tissue are then grown into full plants in accordance with standard techniques. Transgenic seed can be obtained from transgenic flowering plants in accordance with standard techniques. Likewise, non-flowering plants such as potato and sugar beets can be propagated by a variety of known procedures. See, *e.g.*, Newell *et al.* Plant Cell Rep. 10:30-34 (1991) (disclosing potato transformation by stem culture).

In another embodiment of the present invention, the heterologous nucleic acid is transferred from the donor to the genome of the recipient plant via fusion of somatic cells or protoplasts. An advantage of this embodiment is that some of hybridization barriers limiting sexual crossing are bypassed. This technology is more complex than sexual crossing, however, and it can be used only for crosses between species that can be regenerated from a protoplast. Thus, it is preferred to use donor and recipient plants that are unrelated. Examples of such pairings include *Arabidopsis*/cotton, *Arabidopsis*/soybean, *Arabidopsis*/rice and tobacco/soybean. On the other hand, although hybrids between distantly related species (intergeneric, intertribal and interfamilial) have been created using protoplast hybridization, the ability of two phylogenetically distant genomes to cooperate in a hybrid cell has been limited and hybrid cells are often unstable and quickly lose genetic material of one of parental species. See, Gleba & Sytnik, *Monogr. Theor. Appl. Genet.* 8:1-220 (1984), Dudits, *et al.*, *Proc. Nat. Acad. Sci. USA* 84:8434-8438 (1987), and Babiychuk, *et al.*, *Mol. Gen. Genet.* 84:87-91 (1992). Thus, pairings of related donor and recipient plants that are distantly related are more preferred.

The experiments described below are summaries of successful transformation and line conversion for four important crop species (canola, potato, maize and wheat) based on use of specific species hybridization combinations and either transposon- or homologous recombination-based transgene excision/reinsertion. These examples are presented merely to illustrate specific embodiments of the present invention, and are not intended to provide any limitations to the invention not set forth in the claims.

Examples

Example I

Transformation/line conversion of *Brassica* species

10 *Designing the constructs.*

Binary vectors containing components of *Z. mays* transposable element *Spm* within T-DNA borders were made as described below.

Plasmid pIC012 was digested with *Xho*I and *Sma*I, large fragment was gel purified and ligated with *RS* fragment produced by *Xho*I and *Cla*I/Klenow treatment. Resulting plasmid pIC013, containing pNOS-*RS*-3' *OCS* in pUC118, was digested with *Pst*I and *Bcl*I, gel-purified and ligated with large fragment of pIC017 digested with the same enzymes. Plasmid pIC23, containing d*Spm* element with pNOS-*RS*-3' *OCS* was consequently treated with *Hind*III/Klenow and *Bam*H1/Klenow in order to remove *Hind*III and *Bam*H1 sites. Large *Cla*I fragment of pIC23 (-*Bam*H1;-*Hind*III) was cloned into *Cla*I site of pIC201, giving d*Spm* element flanked by p35*S* and *GUS*-3' *NOS* (pIC132). Plasmid pIC132 was digested with *Eco*R1 and *Hind*III. Large fragment was gel-purified and cloned into *Eco*R1 and *Hind*III sites of binary vector based on pRK290 and carrying *NPTII* gene as the plant transformation marker. Resulting plasmid pIC141 was digested with *Ecl*136II and *Hind*III, ligated with 0.3 kb *Xho*I - *Bam*H1 fragment of pIC022 and 7.7 kb *Xho*I-*Sma*I fragment of pIC023 in order to introduce *Spm* transposase under control of 35*S* promoter into the binary vector. Plasmid pIC156 was obtained and used in transformation experiments. Plasmid pIC216 was made in similar way, but *Xho*I - *Bam*H1 fragment of pIC022 was replaced by 0.2 kb *Xho*I-*Bgl*II fragment (p*Spm*) of pIC61.

Cloning steps for another four vectors were only different from the described above by the stages of equipping d*Spm* element with either pNOS:*BAR*-*OCS*3' or with pNOS:*BAR*-*OCS*3' where *BAR* was flanked by two *RS* sites. The plasmid pIC132 was digested with *Pst*I, *Bcl*I, gel-purified and ligated with 1.5 kb *Pst*I, *Bcl*I fragment of pIC016, giving the d*Spm* element with pNOS:*BAR*-*OCS*3'.

For constructs pIC401 and pIC411 large *Xho*I-*Nco*I fragment of pIC01 was ligated with small fragment of *Bsp*HI – *Bam*HI fragment of pIC018 and two *RS* fragments flanked with *Nco*I- *Bsp*HI and *Bgl*II- *Xho*I sites respectively. Resulting plasmid pIC38 consists of *BAR* gene flanked by two *RS* sites. Small *Xho*I fragment of pIC36 was recloned into *Xho*I site of pIC334 giving plasmid pIC342 with pNOS:*RS-BAR-RS-OCS3*'. The last cassette was re-cloned into d*Spm* element as described above.

In summary, constructs obtained consist of transformation marker (NPTII gene conferring resistance to kanamycin), source of *Spm* transposase under control of either 35S or its own promoter, non-autonomous d*Spm* element inserted within p35S:*GUS* excision marker. Three different versions of d*Spm* element were made:

a) d*Spm* contains *NOS* promoter separated by *RS* site (recombination site recognized by R recombinase from *Z. rouxii*) from the terminator of transcription of *OCS* gene. (See: pIC156 and pIC216, Figs. 1 and 2).

b) d*Spm* contains pNOS:*BAR-OCS3*'. (pIC312, pIC31A2, Figs.3 and 4).

c) d*Spm* contains pNOS:*BAR-OCS3*', but *BAR* gene is flanked by two unidirected *RS* sites (pIC401, pIC411, Figs. 5 and 6).

Constructs were tested in *Arabidopsis* using *in planta* transformation procedure. The d*Spm* excision can be easily monitored in primary transformants by the presence of GUS+ sectors after staining of plant tissues with X-gluc. All constructs showed high transposition activity in *Arabidopsis* and were used to obtain and characterize several *Orychophragmus violaceus* transformants.

Line conversion using O. violaceus.

Seed of *Orychophragmus violaceus* were sterilized and germinated *in vitro*. The transformation of *in vitro* grown plants of the species has been done as previously described for *Brassica* species (De Block, *et al.*, *Plant Physiol.*, 91, 694-701 (1989). The constructs used were *Agrobacterium*-based carrying *Spm* transposase along with different versions of non-autonomous d*Spm* element inserted between 35S CaMV promoter and GUS gene (see Fig.1) The plasmids were used to produce transformed *Orychophragmus* plants. Several transgenic plants have been produced and characterized. Two independent transformants containing a single copy insertion have been crossed as male parents to different *Brassica* species (*B.nigra*, *B.juncea*, *B.napus*, *B. carinata*) and *Sinapsis alba* as previously described. In total, approximately 600 crosses were done. The resultant hybrids were allowed to self and the F₁ progeny has been selected for the presence of d*Spm* element

(PCR analysis or phosphinotricin resistance). Those surviving selection were further screened for pure *Brassica* phenotype and for absence of GUS activity, and, finally, tested for absence of either transposase sequences, or species-specific *Orychophragmus* repeats. Finally, co-segregation of dSpm with a *Brassica* chromosome-specific RFLP pattern has been established by analyzing the F₂ progeny.

Line conversion using Arabidopsis thaliana.

In the examples that follow, all experiments were performed as described above except that instead of *O. violaceus*, *A. thaliana* plants were used as male parents. *Arabidopsis* is easy to transform and has short life cycle. These features make *Arabidopsis* excellent candidate for the clipboard species.

Example II

Transformation/line conversion of *Brassica napus*

Seed of *Brassica napus*, var. and those of *Orychophragmus violaceus* were sterilized and germinated *in vitro*. Transformation was performed as described in (De Block, *et al.*, *Plant Physiol.* 91:694-701 (1989). *Orychophragmus* seed was transformed with *Agrobacterium*-based vector pII2 containing gene for R recombinase and a promoterless gene for hygromycin resistance flanked by two *rsx* recombination sites. Rape seed organism was transformed with vector pII3 containing a 35S CaMV promoter with a *RS* recombinant site, so that proper recombination would create an active HPT gene conferring hygromycine resistance. Two independent transformed plants of each species were selected based on molecular analysis of the transgenics. Crosses and analysis of the progeny was performed as in Example II.

Example III

Transformation/line conversion of potato

Experiments were performed as above (Example I) except transgenic *Solanum phureja* was used as a pollen partner. The crosses were performed as described in Hermesen, *et al.*, *Euphytica* 22:244-259 (1973), and primary converted lines were selected as F₀ diploidized dihaploids.

Example IV

Transformation/line conversion of maize

Tripsacum dactyloides line was used in this experiment as a transgene donor. The constructs used were *Agrobacterium*-based as shown in Figure 1, carrying Spm transposase along with non-autonomous dSpm element inserted between 35S CaMV

promoter and GUS gene, the dSpm containing either one *RS* recombination site or one selectable marker (*BAR*) with (pIC401, pIC411) or without (pIC312, pIC31A2) *RS* sites. Transformation of the parental material was essentially performed as described in Hiei, *et al.*, *Plant Mol. Biol.* 35:205-218 (1997). Transgenic plants were crossed with maize, var., and the resultant progeny was selfed. Pure maize-type segregates were screened from among the BC₁ that showed phyosphinotricin resistance or dSpm-specific PCR signal. Those surviving selection were further screened for pure maize phenotype and for absence of GUS activity, and, finally, tested for absence of either transposase sequences, or species-specific *Tripsacum* repeats. Finally, co-segregation of either phosphinotricin resistance or dSpm-specific PCR signal with a maize chromosome-specific RFLP pattern was established by analyzing the BC/F₂ progeny.

Example V

Transformation/line conversion of wheat.

The experiments were performed as in previous example (Example IV) except the crosses were performed as described in Riera-Lizararu & Mujeeb-Kazi, *Crop Sci.* 33:973-976 (1993). Primary converted lines were selected as F₀ diploidized haploids emerging from the crosses.

INDUSTRIAL APPLICABILITY

The present invention is useful in the production of genetically engineered plants that exhibit a wide array of properties that may include enhanced resistance to viruses, fungi, bacterial diseases, pests, pesticides or environmental stress, as well as for the enhancement of other commercially desirable properties such as improved flavor, storage or nutritional properties.

All publications mentioned in this specification are indicative of the level of skill of persons skilled in the art to which this invention pertains. All these publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

Various modifications of the invention described herein will become apparent to those skilled in the art. Such modifications are intended to fall within the scope of the appending claims.

CLAIMS:

1. A method for introducing genetic material into plants, comprising:
preparing a first plant transformed with a heterologous nucleic acid having 5' and 3' excisable flanking sequences that allow movement of said heterologous nucleic acid
5 from one genome to another;
crossing a second plant and the transformed first plant, wherein said first and second plants, upon crossing, produce unstable progeny or demonstrate preferential segregation or sorting out; and
selecting progeny of said second plant of (b) which contain said heterologous
10 nucleic acid.
2. The method of claim 1, wherein said 5' and 3' excisable flanking sequences comprise a transposable element, and wherein said first plant, said second plant or both said first plant and said second plants produce a transposase specific to said transposable element.
3. The method of claim 1, wherein said 5' and 3' excisable flanking sequences are recombination sites and wherein said first plant, said second plant or both said first and second plants produce a recombinase specific to said recombination sites.
4. The method of claim 1, wherein said first plant is *Tripsacum* and said second plant is maize.
5. The method of claim 1, wherein said first plant is *Tripsacum* and said second plant is wheat.
6. The method of claim 1, wherein said first plant is *Tripsacum* and said second plant is barley.
7. The method of claim 1 wherein said first plant is *Tripsacum* and said
25 second plant is oat.
8. The method of claim 1 wherein said first plant is *Orychophragmus* and said second plant is a crucifer.
9. The method of claim 1 wherein said first plant is *Arabidopsis* and said second plant is a crucifer.
10. The method of claim 8 wherein said crucifer is canola.
11. The method of claim 1, wherein said first plant is *Glycine tomentella* and said second plant is soybean.

12. The method of claim 1, wherein said first plant is *Solanum phreja* and said second plant is potato.

13. The method of claim 1, wherein said first plant is maize and said second plant is wheat.

5 14. The method of claim 1, wherein said first plant is maize and said second plant is barley.

15. The method of claim 1, wherein said first plant is maize and said second plant is oats.

10 16. The method of claim 1, wherein said first plant is *Pennisetum* and said second plant is wheat.

17. The method of claim 1, wherein said first plant is *Pennisetum* and said second plant is barley.

18. The method of claim 1, wherein said first plant is *Hordeum bulbosum* and said second plant is barley.

15 19. The method of claim 1, wherein said first plant is *Hordeum bulbosum* and said second plant is wheat.

20. The method of claim 1, wherein said first plant is *Oryza minuta* and said second plant is rice.

20 21. The method of claim 1, wherein said first plant is *Nicotiana dilguta* and said second plant is *Nicotiana tabacum*.

22. The method of claim 1, wherein one or both said first and second plants is cotton carrying a *Se* semigamy mutation.

23. The method of claim 1, wherein one of said first and second plants is soybean carrying a *ms* mutation causing polyembryony.

25 24. The method of Claim 1, wherein said first plant is *Arabidopsis*.

25. A method for introducing genetic material into plants, comprising:

(a) preparing a cell or protoplast of a first plant transformed with a heterologous nucleic acid having 5' and 3' excisable flanking sequences that allow movement of said heterologous nucleic acid from one genome to another;

30 (b) fusing said cell or protoplast with a cell or protoplast of a second plant to produce a fused cell or a fused protoplast, wherein said first and second plants, upon crossing, produce unstable progeny or demonstrate segregation preferential or sorting out;

(c) regenerating whole plants from the fused cell or the fused protoplast; and

-21-

(d) selecting progeny of said regenerated plants of (c) which contain said heterologous nucleic acid.

26. The method of claim 25, wherein said 5' and 3' excisable flanking sequences comprise a recombination site.

5 27. The method of claim 25, wherein said fusing is conducted in medium containing a recombinase specific for said recombination site.

28. The method of claim 25, wherein said first plant species, said second plant species or both said first and second plant species produce a recombinase specific to said recombination sites.

10 29. The method of claim 25, wherein said 5' and 3' excisable flanking sequences comprise a transposable element, and wherein said first plant, said second plant or both said first plant and said second plants produce a transposase specific to said transposable element.

15 30. The method of claim 25 wherein said first plant is *Arabidopsis* and said second plant is cotton.

31. The method of claim 25 wherein said first plant is *Arabidopsis* and said second plant is soybean.

32. The method of claim 25 wherein said first plant is *Arabidopsis* and said second plant is rice.

20 33. A whole plant containing a heterologous nucleic acid, prepared by the method of Claim 1 or Claim 24.

34. Seed or seed parts from the whole plant of Claim 33.

35 Progeny of the plant of Claim 33.

25 36. A fused cell or fused protoplast produced by the method of Claim 25(b).

FIG. 1





FIG. 2



SUBSTITUTE SHEET (RULE 26)




FIG. 4






FIG. 5



204000 24508660

FIG. 6



Please type a plus sign (+) inside this box



PTO/SB/01 (03-01)
Approved for use through 10/31/2002. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION FOR UTILITY OR
DESIGN
PATENT APPLICATION
(37 CFR 1.63)**

☐ Declaration
Submitted
with Initial
Filing

OR

☐ Declaration
Submitted after Initial
Filing (surcharge
(37 CFR 1.16 (e))
required)

Attorney Docket Number

ICON 3.3-001

First Named Inventor

Nikolay V. Kuchuk, et al.

COMPLETE IF KNOWN

Application Number

Filing Date

Group Art Unit

Examiner Name

As a below named inventor, I hereby declare that:

My residence, mailing address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

PROCESS OF RAPID VARIETY-INDEPENDENT PLANT TRANSFORMATION

(Title of the Invention)

the specification of which

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY) May 17, 2000 as United States Application Number or PCT International
Application No. PCT/US00/13555 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or (f), or 365(b) of any foreign application(s) for patent, inventor's or plant breeder's rights certificate(s), or 365 (a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent, inventor's or plant breeder's rights certificate(s), or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

Please type a plus sign (+) inside this box ☐

PTO/SB/01 (03-01)
Approved for use through 10/31/2002. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION — Utility or Design Patent Application

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Customer Number 000530

Direct all correspondence to:		<input checked="" type="checkbox"/> Customer Number or Bar Code Label	000530	OR <input type="checkbox"/> Correspondence address below
Name				
Address				
City		State		ZIP
Country	Telephone		Fax	
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.				
NAME OF SOLE OR FIRST INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor		
Given Name (first and middle [if any])		Family Name or Surname		
Inventor's Signature		Date		
Residence: City	State	Country	Citizenship	
Mailing Address:	Uborevicha Street, 27, Apartment 145			
City	State	ZIP	Country	
NAME OF SECOND INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor		
Given Name (first and middle [if any])		Family Name or Surname		
Inventor's Signature		Date		
Residence: City	State	Country	Citizenship	
Mailing Address:	98 North Park Avenue			
City	State	ZIP	Country	
<input type="checkbox"/> Additional inventors are being named on the _____ supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto.				